

BE for CE-Level Decision Template

Prepared by

Signature: _____/s/ *Diana Perez* _____ *Diana Perez* te: _____ 2/5/2020 _____

Signature: _____/s/ *Shawna Bautista* _____ *Shawna Bautista* date: _____ Feb. 05, 2020 _____

Section I – Screening Tables

Species List Analyzed:

R6 RFSS List Date: ☒ 3/8/2019 (<https://www.fs.fed.us/r6/sfpnw/issssp/agency-policy/>);

☐ - Other Date; Date TEP list acquired from [IPaC](#) Date

Table 1. Species that are not analyzed further

Listed and Proposed Species
The following listed or proposed species or critical habitats are not known or expected to occur in the project area, or are not expected to be affected by the project. As a result, no effect is expected to these listed or proposed species and effects to them are not analyzed further :
All species <input type="checkbox"/>
TEP species considered, but not present : marbled murrelet, northern spotted owl, Oregon spotted frog and its designated critical habitat, gray wolf, North American wolverine, Pacific fisher and its proposed critical habitat, coastal marten
All species except those listed below and analyzed in Section II <input checked="" type="checkbox"/>
The following listed or proposed species or critical habitat is known or expected to occur or may be affected by the project and are analyzed further. Proceed to Section II with these species.
<u>List species here</u> : Southern Oregon/Northern California Coho (SONCC) and its designated critical habitat
Regional Forester's Sensitive Species
The following sensitive species are not known or expected to occur in the project area. As a result, no impact is expected to these sensitive species and impacts to them are not analyzed further :
All species <input type="checkbox"/>
All species except those listed below and analyzed in Section II <input checked="" type="checkbox"/>
The following sensitive species are known or expected to occur in the project area and are analyzed further. Proceed to Section II with these species.
<u>List species here</u> : Tricolored blackbird, Lewis' woodpecker, Purple martin, Pallid bat, Townsend's big-eared bat, Fringed myotis, Oregon branded skipper, Coronis fritillary, Pacific lamprey, KMP steelhead, SONCC Chinook salmon

Section II – Analysis and determination of effect*Table 2. Identification of habitat and analysis of impacts*

Listed and Proposed Species carried forward from Section I		
For each species carried forward from Section I, briefly identify and describe all occupied and unoccupied habitat as it relates to recovery and summarize how the proposed action may directly, indirectly, or cumulatively affect the species or their occupied habitat, or unoccupied habitat required for recovery		
Species	Habitat description	Summary of potential effects from proposed action on species or habitat
SONCC coho	Habitat for SONCC coho salmon is found within the project area (JHSN) in Jackson Creek. However, habitat is not accessible due to a fish passage barrier located outside the project area downstream at the Hanley Rd crossing estimated at river mile 4.3. Distribution and habitat within action area.	SONCC coho and their prey source (invertebrates) as well as macrophytes are not likely to be exposed to pesticides at detectable levels of concern, and thus not likely to be adversely affected by the exposure. Effects are expected to be insignificant or discountable because the amount of pesticides proposed for use that could enter Jackson Creek would be so minor that it cannot be meaningfully evaluated. Use of high risk pesticide (Chlorpyrifos) is restricted to one time use in shadehouses and greenhouses, so coho are unlikely to be exposed. Refer to Fisheries Biological Assessment for more information.
Designated Critical Habitat for SONCC	Designated critical habitat for SONCC coho exists in Jackson Creek on JHSN and upstream to where natural gradients impede migration. The type of habitat for SONCC coho in Jackson Creek at JHSN would be limited to rearing and migration year round. Existing migration barriers and the availability of adequate water flow year round impact the presence of coho in Jackson Creek, refer to the Fisheries Biological Assessment. Critical habitat is not accessible due to a fish passage barrier located outside the project area downstream at the Hanley Rd crossing estimated at river mile 4.3. Critical habitat within action area.	Pesticides proposed for use pose a low risk to fish, amphibians, invertebrates, and macrophytes because they would not reach Jackson Creek at levels that could adversely affect habitat as a result of site-specific conditions and properties. Use of high risk pesticide (Chlorpyrifos) is restricted to one time use in shadehouses and greenhouses, so primary biological features are unlikely to be exposed. Refer to Fisheries Biological Assessment for more information.

Regional Forester's Sensitive Species carried forward from Section I		
For each species carried forward from Section, briefly identify and describe all occupied and unoccupied habitat as it relates to maintaining viability at the Forest Plan scale or preventing a trend towards listing and summarize how the proposed action may directly, indirectly, or cumulatively impact the species or their occupied habitat		
Species	Habitat description	Summary of potential impacts from proposed action on species or habitat
Lewis' Woodpecker	Oregon white oak, ponderosa pine and cottonwood. Reported to winter along Bear Creek near Medford.	No impact to habitat or species. They only winter in project area when pesticides are not used and their food is primarily fruits and acorns.
Purple martin	cropland, hedgerows, grasslands, shrubland, suburban, orchard, and woodlands	No impact to habitat. May impact individuals but will not lead to a trend toward federal listing. Few individuals near project area. Some risk to individuals if consume insects contaminated by oryzalin, but martins are aerial feeders and likelihood of contaminating prey is low.
Tricolored blackbird	Freshwater marshes with cattails, dense willows, Himalayan (Armenian) blackberries	No impact to habitat. May impact individuals but will not lead to a trend toward federal listing. Could possibly forage in nursery fields. Oregon birds represent 1% of total population. Risk to some individuals if they consume enough food contaminated with oryzalin or thiram
Fringed myotis	Breeds in caves, mines, buildings; forested or riparian areas; forages on shrubs and the ground	No impact to habitat or species. Only chlorpyrifos poses risk to insect-eating mammals, and its use is restricted to greenhouse, so prey unlikely to be exposed.
Townsend's big-eared bat	Caves, mines, bridges, rock crevices and old buildings; forages in flight and from foliage	No impact to habitat or species. Only chlorpyrifos poses risk to insect-eating mammals, and its use is restricted to greenhouse, so prey unlikely to be exposed.
Pallid bat	Arid areas, open forests, with rock crevices, caves, old mines, trees or old buildings; forages on ground	No impact to habitat or species. Only chlorpyrifos poses risk to insect-eating mammals, and its use is restricted to greenhouse, so prey unlikely to be exposed.
Species	Habitat description	Summary of potential impacts from proposed action on

Biological Evaluation for Categorical Exclusion

		species or habitat
Oregon branded skipper	420-1500 meters in elevation, hillslopes with flowers, rabbitbrush	No impact to habitat. Azadirachtin, chlorpyrifos, esfenvalerate, and pyrifproxifen pose risk to insects. Only isolated individuals likely to visit nursery. Operational BMP's to reduce exposure. May impact individuals but will not lead to a trend toward federal listing.
Coronis fritillary	Low elevation grasslands, with some shrubs	No impact to habitat. Azadirachtin, chlorpyrifos, esfenvalerate, and pyrifproxifen pose risk to insects. Only isolated individuals likely to visit nursery. Operational BMP's to reduce exposure. May impact individuals but will not lead to a trend toward federal listing.
Pacific Lamprey	Pacific lampreys spawn in habitat similar to that of salmon: gravel bottomed streams at the upstream end of riffle habitat. Adults lay egg nests in some of the same gravel bars as wild salmon and steelhead. Juveniles need areas of low velocity and fine substrates where they burrow, grow and live as filter feeders for 3 to 7 years and feed primarily on diatoms and algae. Lamprey have been found in Jackson Creek, though the exact location is unknown and believed to be in the lower stream reaches of Jackson Creek. Lamprey habitat may exist in Jackson Creek up to a fish passage barrier on river mile 4.3, which is almost 0.5 miles downstream from JHSN. Lamprey and habitat in action area.	Pacific Lamprey and their food source may be exposed to pesticide residues as a result of runoff from JHSN. Residue is not expected to be at detectable levels of concern because the amount of pesticides proposed for use that could plausibly enter Jackson Creek would be insignificant. May impact individuals but will not lead to a trend toward federal listing.
KMP Steelhead	Habitat for steelhead is present in Jackson on JHSN. Historic spawning habitat for summer steelhead is present in Jackson Creek up to and above Jacksonville, Oregon – well past JHSN. Although the upper reaches of Jackson creek around Jacksonville, Oregon dry up in the summer, they can still produce summer steelhead. In 2019 steelhead fry were observed upstream of the three way stop of Main Street and Jacksonville Highway. Steelhead and habitat present in action area.	KMP Steelhead and their food source may be exposed to pesticide residues as a result of runoff from JHSN. Residue is not expected to be at detectable levels of concern because the amount of pesticides proposed for use that could plausibly enter Jackson Creek would be insignificant. May impact individuals but will not lead to a trend toward federal listing.
Species	Habitat description	Summary of potential impacts from proposed action on

Biological Evaluation for Categorical Exclusion

		species or habitat
SONCC Chinook salmon	Fall run Chinook Salmon is limited to the first 1.5 river miles of Jackson Creek, basically up to the confluence of Jackson Creek and Dean Creek, which is significantly downstream of JHSN (a little under 5 miles). Spawning habitats are characterized by large cobbles and sufficient flows to facilitate oxygen delivery to developing embryos. SONCC chinook observance below action area.	No impact to habitat or species. Distribution outside of action area.

Table 3. Determination of effect

Listed and Proposed Species and their Critical Habitat carried forward from Section I	
The proposed action:	Species
Will have no effect (qualifies for CE)	(All terrestrial listed species are no effect; none present in project area)
<u>Proposed</u> - Not Likely to Jeopardize the Continued Existence Of The Species Or Result In Destruction Or Adverse Modification Of Proposed Critical Habitat*. <u>Listed</u> - May affect, but not likely to adversely affect* (qualifies for CE)	SONC coho and its designated critical habitat
<u>Proposed</u> - Likely to Jeopardize the Continued Existence Of The Species Or Result In Destruction Or Adverse Modification Of Proposed Critical Habitat*. <u>Listed</u> - May affect, and is likely to adversely affect* (See Attachment - 1 to the Chief's Letter Dated October 12, 2018, Pertaining to Significance under the Endangered Species Act and National Environmental Policy Act to determine if your project qualifies for CE)	
Regional Forester's Sensitive Species carried forward from Section I	
The proposed action:	Species
Will have no impact (qualifies for CE)	Lewis' woodpecker, , Townsend's big-eared bat, Pallid bat, SONCC chinook
Will have a beneficial impact (qualifies for CE)	Indirectly, all species, as native plants grown at the nursery are used for reforestation and restoration of habitat.

Regional Forester's Sensitive Species carried forward from Section I	
May Impact Individuals Or Habitat, But Will Not Likely Contribute To A Trend Towards Federal Listing or Cause A Loss Of Viability To The Population Or Species (qualifies for CE)	Purple martin, Tricolored blackbird, Fringed myotis, Oregon branded skipper, Coronis fritillary, Pacific Lamprey, KMP Steelhead
Will Impact Individuals Or Habitat With A Consequence That The Action May Contribute To A Trend Towards Federal Listing Or Cause A Loss Of Viability To The Population or Species (likely does NOT qualify for CE)	

* Biological Assessments will need to be completed and consultation will need to be completed prior to signing the decision for these determinations.

Section III –

Project Design Criteria and Best Management Practices for dealing with adverse effects

Project Design Criteria and Best Management Practices to be used to reduce negative effects/impacts of the project aimed at helping achieve, maintain, or restore project eligibility for CE – These PDCs and BMPs should be collaboratively developed, ideally during the Plan to Project phase, with the responsible official and other specialists. If these are not followed, it may change the determination of effects above.

Table 4. Project Design Criteria and Best Management Practices for removing, avoiding, or compensating for any adverse effects and notes for particular species

Species	PDCs and BMPs
Fringed myotis, Townsend's big-eared bat, pallid bat, Oregon branded skipper, Coronis fritillary	Chlorpyrifos and pyriproxyfen are limited to use in greenhouses and shadehouses only, making exposure unlikely.
Oregon branded skipper, Coronis fritillary	Insecticide use is discontinued on native forbs when flowering begins.
SONCC coho and its designated critical habitat, Pacific Lamprey, KMP Steelhead, SONCC Chinook	<p>The design features are based on site-specific pesticide risk assessments and their possible effects to aquatic species.</p> <ul style="list-style-type: none"> ✓ Minimize overall use of pesticides by implementing integrated pest management methods; continue to utilize mechanical, manual, biological and cultural methods where cost-effective. ✓ Minimize use of chemicals that are more mobile and persistent (oxyfluorfen, propiconazole), especially prior to or during the rainy season. ✓ Minimize use of pesticides that have been detected with increasing

	<p>frequency in local aquatic habitats (glyphosate, oxyfluorfen).</p> <ul style="list-style-type: none">✓ Use chlorpyrifos only in the greenhouses, limited to one application per greenhouse per year. Favor other effective products.✓ Manage timing of spraying and irrigation to minimize run off into Jackson Creek.✓ Use cover crops and reduce tillage where appropriate.✓ Reduce potential for drift by only broadcast spraying when wind speed is between 2 and 8 miles per hour, using the largest feasible nozzle size and lowest spray height, and protecting non-target resources with drift shields. Avoid broadcast within 100 feet of flowering plants when pollinators are present.✓ Promptly clean up all spills, including treated seed. (Please note no reportable spills of pesticide have occurred at the nursery.)✓ Err on the side of caution when interpreting label guidance.✓ Use adjuvants approved for use on aquatic environments by Washington state (refer to Fisheries Biological Assessment).
--	---

Section IV – References

- Beedy, E. C., W. J. Hamilton, III, R. J. Meese, D. A. Airola, and P. Pyle (2018). Tricolored Blackbird (*Agelaius tricolor*), version 3.1. In *The Birds of North America* (P. G. Rodewald, Editor). Cornell Lab of Ornithology, Ithaca, NY, USA. <https://doi.org/10.2173/bna.tribla.03.1>
- Bird Studies Canada and NABCI. 2014. BirdConservation Regions. Published by Bird Studies Canada on behalf of the North American Bird Conservation Initiative. <http://www.birdscanada.org/research/gislab/index.jsp?targetpg=bcr> Accessed: July 18, 2019
- Blevins, E. 2016. Species Fact Sheet: *Hesperia colorado oregonia* (W. H. Edwards, 1883). USDA Forest Service, Pacific Northwest Region. Portland, OR. Available online at: <https://www.fs.fed.us/r6/sfpnw/issssp/species-index/fauna-invertebrates.shtml> Accessed: 25 June 2019
- Bureau of Land Management, Medford District, Ashland Resource Area. August 2001. West Bear Creek Watershed Analysis. Version 1.1. https://www.blm.gov/or/districts/medford/plans/files/WA/WestBearCk_WA_acc.pdf
- Cameron, S. A., J. D. Lozier, J. P. Strange, J. B. Koch, N. Cordes, L. F. Solter, and T. L. Griswold. 2011a. Patterns of widespread decline in North American bumble bees. *Proceedings of the National Academy of Sciences* 108:662–667.
- Christensen, K.; Harper, B.; Luukinen, B.; Buhl, K.; Stone, D. 2009. Chlorpyrifos Technical Fact Sheet; National Pesticide Information Center, Oregon State University Extension Services. <http://npic.orst.edu/factsheets/archive/chlorptech.html>
- Clayton, D. Personal communication and Personal observation. 2015. Dave Clayton is the Wildlife Program Manager, Rogue River-Siskiyou National Forest, Medford, Oregon.
- COSEWIC (Committee on the Status of Endangered Wildlife in Canada). 2013. COSEWIC assessment and status report on the Oregon Branded Skipper *Hesperia colorado oregonia* in Canada. Committee on the Status of Endangered Wildlife in Canada. Ottawa. ix + 51 pp.

- (<https://www.canada.ca/en/environment-climate-change/services/species-risk-public-registry.html>).
- Cloyd, R.A. 2015. Explaining azadirachtin and neem. Greenhouse Product News, November 2015. <https://gpnmag.com/article/explaining-azadirachtin-and-neem/> Accessed: June 14, 2019.
- Cross, S.P., H. Lauchstedt, and M. Blankenship. 1997. Bat studies in the Ashland and Applegate Districts of the Rogue River National Forest. Final Report. Southern Oregon University.
- Cross, S. P., and D. Waldien. 1995. Survey of bats and their habitats in the Roseburg District of the BLM in 1994. Final Report. Southern Oregon State College, Ashland, Oregon.
- Csuti, B.; T.A. O’Neil; M.M. Shaughnessy; E.P. Gaines; J.C. Hak. 2001. Atlas of Oregon Wildlife: Distribution, Habitat, and Natural History. Oregon State University Press. Corvallis, OR 524 pp.
- EPA (Environmental Protection Agency). 1998. Reregistration eligibility decision (RED) *Bacillus thuringiensis*. Prevention, Pesticide and Toxic Substances. Washington, D.C. 170 pp.
- EPA (Environmental Protection Agency). 2002. Biopesticide registration action document: sodium carbonate peroxyhydrate, PC Code 128860. Biopesticides and Pollution Prevention Division, Office of Pesticide Programs. Washington, D.C. 31pp.
- EPA (Environmental Protection Agency). 2004. Environmental fate and ecological risk assessment for the reregistration of thiram. Environmental Fate and Effects Division, Office of Pesticide Programs. Washington, D.C. 71pp.
- EPA (Environmental Protection Agency). 2006. Ecological risk assessment for the reregistration of propiconazole (revised). Office of Prevention, Pesticides, and Toxic Substances. Washington, D.C. 206pp.
- EPA (Environmental Protection Agency). 2015. Reregistration eligibility decision (RED): Metalaxyl. Office of Prevention, Pesticides, and Toxic Substances. Washington, D.C. 335 pp.
- EPA (Environmental Protection Agency). 2012. Biopesticides registration document. Cold pressed neem oil, PC Code 025006. Office of Pesticide Program, Biopesticides and Pollution Prevention Division. Washington, D.C. 21pp.
- EPA (Environmental Protection Agency). 2017. Preliminary ecological risk assessment for the registration review: pyriproxyfen. Environmental Fate and Effects Division, Office of Pesticide Programs. Washington, D.C. 102pp.
- Freeman, M. 2019. A gourd to call home. Mail Tribune, Friday, March 22nd. <https://mailtribune.com/news/videos/gourds-give-rare-swallows-a-place-to-call-home>
- Fürst MA, McMahon DP, Osborne JL, Paxton RJ, Brown MJF. 2014. Disease associations between honeybees and bumblebees as a threat to wild pollinators. *Nature* 506:364–366.
- Galen, C. 2003. Lewis’ woodpecker. Pp. 351-352 in *Birds of Oregon: A General Reference*. D.B. Marshall, M.G. Hunter, and A.L. Contreras, Eds. Oregon State University Press, Corvallis, OR.
- Gervais, J. 2016. Conservation Assessment for the Pallid Bat (*Antrozous pallidus*) in Oregon and Washington. Unpub. Report prepared for USDA Forest Service, Region 6 and USDI Bureau of Land Management, Oregon and Washington. 44 pp. Available at: <https://www.fs.fed.us/r6/sfpnw/issssp/species-index/fauna-mammals-bats.shtml> Accessed: 30 July 2019.

- Gervais, J. 2017. Conservation Assessment for the Townsend's big-eared bat (*Corynorhinus townsendii*) in Oregon and Washington. Unpub. Report prepared for USDA Forest Service, Region 6 and USDI Bureau of Land Management, Oregon and Washington. 47 pp. Available at: <https://www.fs.fed.us/r6/sfpnw/issssp/species-index/fauna-mammals-bats.shtml> Accessed: 30 July 2019.
- Gervais, J. 2017. Conservation Assessment for the fringed myotis (*Myotis thysanodes*) in Oregon and Washington. Unpub. Report prepared for USDA Forest Service, Region 6 and USDI Bureau of Land Management, Oregon and Washington. 47 pp. Available at: <https://www.fs.fed.us/r6/sfpnw/issssp/species-index/fauna-mammals-bats.shtml> Accessed: 30 July 2019.
- Goulson D. 2010. Bumblebees: behaviour, ecology, and conservation. Oxford University Press.
- Hallman, A.; Bond, C.; Buhl, K.; Stone, D. 2015. Pyriproxyfen General Fact Sheet; National Pesticide Information Center, Oregon State University Extension Services. <http://npic.orst.edu/factsheets/pyriprogen.html>
- Hatfield R, Jepsen S, Mader E, Black SH, Shepherd M. 2012. Conserving Bumble Bees. Guide-lines for Creating and Managing Habitat for America's Declining Pollinators. Available from http://www.xerces.org/wp-content/uploads/2012/06/conserving_bb.pdf (accessed 25 June 2019). Hatfield RG, Colla SR, Jepsen S, Richardson LL, Thorp RW, Foltz-Jordan S. 2014a. IUCN Assessments for North American *Bombus* spp. for the North American IUCN Bumble Bee Specialist Group. The Xerces Society for Invertebrate Conservation, Portland, OR.
- Hatfield, R., Jepsen, S., Thorp, R., Richardson, L. & Colla, S. 2015. *Bombus suckleyi*. The IUCN Red List of Threatened Species 2015: e.T44937699A46440241. <http://dx.doi.org/10.2305/IUCN.UK.2015-2.RLTS.T44937699A46440241.en>. Downloaded on 25 June 2019.
- Hatfield, R., C. Fallon, and M. Blackburn. 2018. Mardon skipper (*Polites mardon*) distance sampling surveys at four sentinel sites in Oregon and Washington: Year 5. Status report to the U.S. Forest Service, Bureau of Land Management, and the interagency special status/sensitive species program (ISSSP). Unpub. Report. 31 pp. Available at: <https://www.fs.fed.us/r6/sfpnw/issssp/species-index/fauna-invertebrates.shtml> Accessed: 25 June 2019.
- Hopwood J., M. Vaughan, M. Shepherd, E. Mader, and S. H. Black. 2012. Are Neonicotinoids Killing Bees? A review of research into the effects of neonicotinoid insecticides on bees, with recommendation for action. Tech. rep., The Xerces Society. <http://www.xerces.org/neonicotinoids-and-bees/>
- Horvath, E.G. 1999. Oregon purple martins. <http://www.purple-martin.org/WesternMartins/OregonHorvath.htm>. Accessed: 25 June 2019.
- Horvath, E.G. 2003. Purple martin. Pp. 428-430 in *Birds of Oregon: A General Reference*. D.B. Marshall, M.G. Hunter, and A.L. Contreras, Eds. Oregon State University Press, Corvallis, OR
- IPCS (International Programme on Chemical Safety). 1999 Environmental Health Criteria, No 217. *Bacillus thuringiensis*, 1999, 105pp. available online at: <http://www.inchem.org/documents/ehc/ehc/ehc217.htm#SectionNumber:1.5>
- Janes, S. 2017. Purple martins are moving in. Mail Tribune, Friday, August 25th.

- Jordan, S.F. 2011. Species Fact Sheet: *Speyeria coronis* nr. *coronis* (Behr, 1864). Available at: <https://www.fs.fed.us/r6/sfpnw/issssp/species-index/fauna-invertebrates.shtml> Accessed: 25 June 2019.
- Jordan, S.F. 2012. Species Fact Sheet: *Hesperia colorado oregonia* (W. H. Edwards, 1883). Available at: <https://www.fs.fed.us/r6/sfpnw/issssp/species-index/fauna-invertebrates.shtml> Accessed: 25 June 2019.
- Justin, John. 2019. J. Herbert Stone Nursery Manager. USDA Forest Service, Region 6. Personal communication.
- Mason, Bruce, and Girard, Inc. (MB&G). 2001. Biological Assessment: Highway 62 Corridor Solutions Project, North Medford Interchange. Prepared for the Oregon Department of Transportation.
- Mitchell, C., J. Doerr, S. Colyer, and R. Espinosa. 2016. 2015 SW Oregon Integrated Western Bumble Bee Survey Project: Summary Report of Findings. Unpub. Report. USDA Forest Service. Available at: <https://www.fs.fed.us/r6/sfpnw/issssp/species-index/fauna-invertebrates.shtml>. Accessed: 25 June 2019.
- NMFS (National Marine Fisheries Service). 2007. 2007 Federal Recovery Outline, Southern Oregon/Northern California Coast Coho Salmon. National Marine Fisheries Service. Southwest Region. Arcata Area Office. Arcata, California. December 12, 2007.
- NMFS (National Marine Fisheries Service). 2014. Final Recovery Plan for the Southern Oregon/Northern California Coast Evolutionarily Significant Unit of Coho Salmon (*Oncorhynchus kisutch*). National Marine Fisheries Service. Arcata, CA.
- NYDEC (New York Department of environmental Conservation). 2015. Active ingredient data package: metalaxyl and mefenoxam. Bureau of Pest Management, Pesticide Product Registration Section. 72 pp.
- Norris, L.A.; Lorz, H.W.; Gregory, S.V. 1991. Forest chemicals. American Fisheries Society Special Publication. 19:207-296.
- ODFW (Oregon Dept. of Fish & Wildlife). 2019. Restoration opportunities on Jackson Creek. Report by Daniel Van Dyke, ODFW Rogue District fish Biologist. June 2019.
- ODFW (Oregon Dept. of Fish & Wildlife). 2019. Wildlife viewing: purple martin. <https://myodfw.com/wildlife-viewing/species/purple-martin> Accessed: December 30, 2019.
- Oregon Flora Project. 2019. Atlas. Roemer's fescue map. Oregon Flora Project, Oregon State Univ., Corvallis, OR. <http://www.oregonflora.org/atlas.php> Accessed: December 30, 2019.
- Pesticide Stewardship Partnership. 2015-17 Biennial Summary. Middle Rogue. Produced by Oregon Water Quality Pesticide Management Team. <https://www.oregon.gov/ODA/shared/Documents/Publications/PesticidesPARC/MiddleRogueSummary.pdf>
- Pierson, E. D., M. C. Wackenhut, J. S. Altenbach, P. Bradley, P. Call, D. L. Genter, C. E. Harris, B. L. Keller, B. Lengus, L. Lewis, B. Luce, K. W. Navo, J. M. Perkins, S. Smith, and L. Welch. 1999. Species conservation assessment and strategy for Townsend's big-eared bat (*Corynorhinus townsendii townsendii* and *Corynorhinus townsendii pallescens*). Idaho Conservation Effort, Idaho Department of Fish and Game, Boise, Idaho.

- Purple Martin Conservation Association (PMCA). 2006. The Purple Martin Conservation Association website, in cooperation with Edenborough University of Pennsylvania. <http://www.purplemartin.org/> Accessed: December 30, 2019.
- Pyle, R.M. 2002. Butterflies of Cascadia. Seattle Audubon Society, Seattle, Washington. 420 pp.
- Reilly, J. and S.H. Black. 2011. Survey Results for the Coronis Fritillary in Southwest Oregon. Unpublished Survey Report. Available online at: <https://www.fs.fed.us/r6/sfpnw/issssp/species-index/fauna-invertebrates.shtml>. Accessed: 25 June 2019.
- Reilly, J. 2019. Personal communication. Medford District, BLM, Biologist. Phone call with Shawna Bautista, July 30, 2019.
- Rockwell, S. 2019. Conservation assessment for purple martin (*Progne subis*). Unpub. Report for USDA Forest Service and USDI Bureau of Land Management Interagency Special Status and Sensitive Species Program, Portland, OR. 73 pp.
- Rogue Valley Council of Governments. 2001. Bear Creek Watershed Assessment. Phase II – Bear Creek Tributary Assessment. <https://www.worldcat.org/title/bear-creek-watershed-assessment-phase-ii-bear-creek-tributary-assessment/oclc/50223268>
- SERA (Syracuse Environmental Research Associates). 2004. Clopyralid Human Health and Ecological Risk Assessment. Report produced for USDA Forest Service, Arlington, Virginia. 154pp.
- SERA (Syracuse Environmental Research Associates). 2004. Dicamba Human Health and Ecological Risk Assessment. Report produced for USDA Forest Service, Arlington, Virginia. 179 pp.
- SERA (Syracuse Environmental Research Associates). 2004. Imazapic Human Health and Ecological Risk Assessment Final Report. Report produced for USDA Forest Service, Arlington, Virginia. 110 pp.
- SERA (Syracuse Environmental Research Associates). 2006. Oxyfluorfen Human Health and Ecological Risk Assessment. Report produced for USDA Forest Service, Arlington, Virginia. 229 pp.
- SERA (Syracuse Environmental Research Associates). 2006. 2,4-D Human Health and Ecological Risk Assessment. Report produced for USDA Forest Service, Arlington, Virginia. 245 pp.
- SERA (Syracuse Environmental Research Associates). 2007. Aminopyralid Human Health and Ecological Risk Assessment – Final Report. Report produced for USDA Forest Service and National Park Service, Atlanta, Georgia. 231pp.
- SERA (Syracuse Environmental Research Associates). 2011. Glyphosate Human Health and Ecological Risk Assessment Final Report. Report produced for USDA Forest Service, Atlanta, Georgia. 336 pp.
- SERA (Syracuse Environmental Research Associates). 2004. Imazapyr Human Health and Ecological Risk Assessment Final Report. Report produced for USDA Forest Service, Atlanta, Georgia. 215 pp.
- SERA (Syracuse Environmental Research Associates). 2014. Dazomet Soil Incorporation: WorksheetMaker Workbook Documentation Final Report. Report submitted to USDA Forest Service, Morgantown, W. Virginia. 30 pp.

- SERA (Syracuse Environmental Research Associates). 2015. Chlorothalonil: WorksheetMaker Workbook Documentation Final Report. Report submitted to USDA Forest Service, Morgantown, W. Virginia. 38 pp.
- SERA (Syracuse Environmental Research Associates). 2015. Chlorpyrifos: WorksheetMaker Workbook Documentation Final Report. Report submitted to USDA Forest Service, Morgantown, W. Virginia. 33 pp.
- SERA (Syracuse Environmental Research Associates). 2015. Mancozeb: WorksheetMaker Workbook Documentation Final Report. Report submitted to USDA Forest Service, Morgantown, W. Virginia. 43 pp.
- SERA (Syracuse Environmental Research Associates). 2015. Oryzalin: WorksheetMaker Workbook Documentation Final Report. Report submitted to USDA Forest Service, Morgantown, W. Virginia. 32 pp.
- Shapiro and Associates, Inc., 2001. Biological Assessment for Coho Salmon and Steelhead Under National Marine Fisheries Service Jurisdiction for the Proposed Replacement of the Bear Creek (Kirtland Rd) Bridge, Jackson County, Oregon. Prepared for Sverdrup Civil, Inc., Lake Oswego, Oregon.
- Snider, R. and S. Godwin. 2016. 2016 Western Bumble Bee Surveys: Medford Bureau of Land Management, Ashland Resource Area. Unpub. Report. Medford, OR 10pp. Available online at: <https://www.fs.fed.us/r6/sfpnw/issssp/species-index/fauna-invertebrates.shtml>
- Spencer, K. 2003. Pp. 578-580 *in* Birds of Oregon: A General Reference. D.B. Marshall, M.G. Hunter, and A.L. Contreras, Eds. Oregon State University Press, Corvallis, OR.
- Tetrahedron, Inc. 2017. Pendimethalin Human Health and Ecological Risk Assessment Final Report-corrected. Report produced for USDA Forest Service, Morgantown, W. Virginia. 41 pp.
- Tetrahedron, Inc. 2017. Prodiamine Human Health and Ecological Risk Assessment Final Report-corrected. Report produced for USDA Forest Service, Morgantown, W. Virginia. 47 pp.
- Tetrahedron, Inc. 2018. Esfenvalerate Human Health and Ecological Risk Assessment Final Report. Report produced for USDA Forest Service, Morgantown, W. Virginia. 49 pp.
- Tetrahedron, Inc. 2018. Iprodione Human Health and Ecological Risk Assessment Final Report-corrected. Report produced for USDA Forest Service, Morgantown, W. Virginia. 44 pp.
- Tetrahedron, Inc. 2018. Thiophanate-methyl Human Health and Ecological Risk Assessment Final Report. Report produced for USDA Forest Service, Morgantown, W. Virginia. 62 pp.
- Thorp, R. W., D. S. Horning and L. L. Dunning. 1983. Bumble bees and cuckoo bumble bees of California (Hymenoptera: Apidae). Bulletin of the California Insect Survey 23: viii.
- Tierney, K.B., Baldwin, D.H., Hara, T.J., Ross, P.S., Scholz, N.L., and Kennedy, C.J. 2010. Olfactory toxicity in fishes. Aquatic Toxicology, 96:2-26.
- Tricolored Blackbird Working Group. 2007. Conservation Plan for the Tricolored Blackbird (*Agelaius tricolor*). Susan Kester (ed.). Sustainable Conservation. San Francisco, CA. 55pp.
- Turner, Larry, PhD. March 31, 2003. Chlorpyrifos Analysis of Risks to Endangered and Threatened Salmon and Steelhead. US EPA, Environmental Field Branch, Office of Pesticide Programs.

- U.S. Environmental Protection Agency. July 20, 1984. EEB Chemical Profile: Oxyfluorfen. US EPA, Washington, DC.
- U.S. Fish and Wildlife Service. 2008. Birds of Conservation Concern 2008. United States Department of Interior, Fish and Wildlife Service, Division of Migratory Bird Management, Arlington, Virginia. 85 pp. [Online version available at: <http://www.fws.gov/migratorybirds/>]
- Vogt, J. 2004. Upper Rogue Smolt Trapping Project, 2001. Oregon Department of Fish and Wildlife Rogue Gish District Report. August 2004.
<http://rainbow.dfw.state.or.us/nrimp/information/smoltreports.htm>
- Warren, A.D. 2005. Butterflies of Oregon: their taxonomy, distribution, and biology. Lepidoptera of North America 6. Contributions of the C.P. Gillette Museum of Arthropod Diversity. Colorado State University, Fort Collins, Colorado. 408 pp.
- Wiggins, D. (2005, March 31). Purple Martin (*Progne subis*): a technical conservation assessment. [Online]. USDA Forest Service, Rocky Mountain Region. Available: <http://www.fs.fed.us/r2/projects/scp/assessments/purplemartin.pdf> [date of access].
- Williams P, Colla S, Xie Z. 2009. Bumblebee Vulnerability: Common Correlates of Winners and Losers across Three Continents. *Conservation biology: the journal of the Society for Conservation Biology* 23:931–940.
- Williams PH, Osborne JL. 2009. Bumblebee vulnerability and conservation world-wide. *Apidologie* 40:367–387.
- Williams, P. H., R. W. Thorp, L. L. Richardson, and S. Colla. 2014. *Bumble Bees of North America: An Identification Guide*. Princeton University Press. 208 pp.
- Williams, P. H., R. W. Thorp, L. L. Richardson, and S. Colla. 2014. *Guide to the Bumble Bees of North America*. Princeton University Press.
- WHO (World Health Organization). 2012. WHO Specifications and Evaluations for Public Health Pesticides; *Bacillus thuringiensis* ssp *israelensis* strain AM65-52. World Health Organization. 52 pp.
- Xerces Society. 2012. Database of records from Bumble Bee Citizen Monitoring Project (2008-2012). Maintained by Rich Hatfield, Xerces Society.